

Publish your terrestrial model data with ESS-DIVE

Charuleka Varadharajan

ESS-DIVE Deputy Lead, Lead: Community Engagement













Objectives



- What are model data?
- ESS-DIVE's community-oriented process for model-data archiving
- New guidelines for storing model data with ESS-DIVE
- Other ESS-DIVE capabilities for hosting model data
- Feedback on additional needs for publishing model data on ESS-DIVE.

Learn about ESS-DIVE's new guidelines for archiving model data and apply it to your datasets!

What does the "model data" mean to you?



What are "model data"?




- Model outputs: Various dimensions and resolutions of final raw output files, spin-up output files, restart files, test data files, higher level output files corresponding to figures or tables
- Model input files: e.g. forcings, parameters
- Metadata files
- Model code
- Scripts for model set-up and initialization; parameterization; post-processing; and visualizations.
- Visualization files

Model data on ESS-DIVE

Walker A; De Kauwe M; Medlyn B; Zaehle S; Asao S; Dietze M; El-Masri B; Hanson






Terrain



Now available: 34 years (!) of high-resolution (1km) WRF dynamically downscaled meteorological data for the East and Taylorriver watersheds (CO). data.ess-dive.lbl.gov/datasets/doi:1.... This took many cpu-hours to produce (thanks @INL). Below: Annual precipitation (mm) in the East



8:53 AM · Feb 18, 2022 · Twitter Web App

Rudisill et al:

https://data.ess-dive.lbl.gov/vie w/doi:10.15485/1845448



Process to develop guidelines for terrestrial model data archiving



Review of data systems that store model data

		Provides data contr	ibutor guideline
Data center	Storage limit per data publication	Model-data specific?	Other?
National Science Foundation Arctic Data Center	No limit	Yes	Yes
Oak Ridge National Laboratory DAAC	NA^1	Yes	Yes
NASA's Earth Observing System Data and Information System (EOSDIS)	NA ¹	NA^1	Yes
U.S. DOE ESS-DIVE	$10GB/500\ GB^2$	No	Yes
Dryad	300 GB ²	No	Yes
Zenodo	50 GB	No	No
Earth System Grid Federation (ESGF)	NA^1	NA^1	NA^1

¹NA: Not available, i.e. no public information found.

²Limit on size of individual files. For ESS-DIVE, 10GB is the default file size limit, and can be increased upto 500GB by request. Files >500GB are considered upon review.





NSF Earthcube Model Data Rubric/ AGU guidelines

EARTHCUBE								
		termining what simulation output:				inicate knowledge.		
		by a combination of the simulation accessed at: https://modeldatarch						
tubric usage instructions an	d Use Case Examples can be	accessed at: https://modeldatarch	.gitnub.io/rubrics-worksr	eets/Rubric-instruction	is-and-use-Cases.pdr			
	Simulation / Expe	riment Descriptors	Simulation	/ Experiment Descrip	otor Classes			
			Preserve Less Output		Preserve More Output			
Big Picture Question	Descriptor	Descriptor definition	Class 1	Class 2		User Entered Score. (Integers only) Score Range: 1 -Class 1 2 -Class 2 3 -Class 3	Suggested Weight (If score > 1)	Weighted Score (Weighted Score = Entered Score x Recommended Weighting when score > 1)
Section Theme: Simulation	Workflow Accessibility			Model source code is				
	Model Source Code Availability	How accessible is this particular version of the model/code? Are there IP barriers, embargo periods for new model development?	Community validated wersion of a highly accessible model was used.	shareable, but specific changes were implemented that make it unique. Code is lightly documented.	Model source code is difficult to acquire		.1	
Would it be straightforward for others in your academic discipline to reru your simulation model run workflow steps?	Model Source Code Documentation/Ease of use	is the source code well documented and easy to use?	Source code is well documented and easy to install and run.		There is very little code and supporting documentation. Source code is difficult to understand and manage.		1	
	Model Compute Platform/System Dependencies	How specialized of a platform is needed to execute the model (specific hardware, compilers, software libraries needed)?	Does not require special hardware, niche software libraries, and licensed compilers to execute. This could include a containerized version of a model.		Requires resources that are more difficult to get access to. E.g. specialized HPC, niche software libraries, and licensed compilers.		.1	
	Simulation Input Accessibility	How much effort is it to get and manage all the inputs used by the simulation?	Simulation inputs/boundary conditions are easy to acquire & manage.		If simulation inputs/boundary conditions are difficult to acquire & manage, retaining output lowers burden for others under to re-run model or use outputs.		1	
					Section Total Raw Score. (Min=4, Max=12)		Section Total Weighted Score. (Min=4, Max=12)	
Section Theme: Simulation	Post Processing Workflow	Accessibility						
Thene. Sululano	Post Processing Source Code Availability	How accessible is this particular version of the post processing code? Are there IP barriers, embargo periods for new model development?	Community validated wersion of a highly accessible post processing workflow was used.	Post processing source code is shareable, but specific changes were implemented that make it unique. Code is lightly documented.	Post processing source code is difficult to acquire		1	
Would it be straightforward for others in your academic	Post Processing Source Code Documentation/Ease of use	is the post processing source code well documented and easy to use?	Source code is well documented and easy to install and run.		There is very little code and supporting documentation. Source code is difficult to understand and manage.		1	

Community Feedback

Community

Needs

Final Guidelines and paper



Background Research




Case studies with published model data



Final Whitepaper



Draft White Paper



Community

Consensus

Questions to Modelers

ECC-DIVE

- Gathered information on
 - Models used
 - File storage and specifications
 - What is worth archiving and how long is the data useful?
 - Archiving protocols
 - Features needed from ESS-DIVE
- Circulated widely to ESS community and conducted 2 webinars on this feedback

1.1 Model data background and storage needs:

- 1. Name the model(s) you would like to archive data from:
- 2. How are your data represented spatially? (e.g., 1 km² resolution):
- 3. Temporal discretization and range of dataset (e.g., hourly time-step for 100 years):
- 4. Number of files for a typical simulation to be archived:
- 5. Average file size for a typical simulation to be archived:
- 6. Types of file formats for a typical simulation to be archived (e.g., netCDF):

1.2 What's worth archiving, why, and for how long?

- Which model data are worth archiving? (Rate each of the following on a scale from 1-5 as Not important at all, Not so important, Somewhat important, Very important, or Extremely important)
 - a. Model inputs
 - b. Model outputs
 - c. Metadata
 - d. Model code
 - e. Other scripts
 - f. Model testing data
- g. Other: Describe any other type of model data worth archiving
- 2. In general how long do you think model data remains useful? (multiple choice)
 - a. <1 year
 - b. 1-2 years
 - c. 5-10 years
 - d. <10 years
- Rate the importance of having these features in a model data repository on a scale of 1-5 (Not important at all, Not so important, Somewhat important, Very important, or Extremely important)
 - a. Sharing of data
 - b. Data preservation (for time period indicated above)
 - c. Complete model data packages that can reproduce the model outputs
 - d. Clear documentation
 - e. Usability of archived model data
- Describe any additional considerations that are important to informing the development of a successful
 model data archive.

1.3 Approaches to archiving model data

- 1. Does your group currently archive model data? (yes or no)
- 2. Describe your group's approach to archiving model data.
- 3. Would you be willing to share documentation of your model data archiving protocol?
- 4. How satisfied are you with your model data archive? (0: Very dissatisfied to 5: Very satisfied)

1.4 Last thoughts

- 1. Note any recommendations you have for model data storage options.
- 2. Would your group be willing to learn a new method for data archiving? (yes, no, absolutely not, maybe)
- 3. Note any other thoughts or comments here.



Participants and projects involved

Researcher	DOE Project(s) Represented
Charlie Koven	NGEE-Tropics, NGEE-Arctic
Jitu Kumar	NGEE-Tropics, NGEE-Arctic
Dipankar Dwivedi	Watershed Function SFA
Anthony Walker	NGEE-Tropics, FACE-MDS, Oakridge TES SFA, RUBISCO, NGEE-Arctic
Xingyuan Chen	PNNL SBR SFA, IDEAS, EXOSHEDS
Scott Painter	NGEE-Arctic, IDEAS, Oakridge Mercury SFA, Exosheds
Dan Ricciuto	Oakridge TES SFA
Qing Zhu	E3SM, RUBISCO
Ethan Coon	NGEE-Arctic, Exasheds
Maoyi Hung	NGEE Tropics, SBC SFA
Kate Maher	SLAC SFA
Ahmad Jan	NGEE-Arctic

Models used across DOE projects



- ELM
- ELM-FATES
- ELM-BeTR
- CLM
- PFLOTRAN
- CABLE
- SDVGM
- GDAY
- ED2
- LPJ-GUESS

- MAAT
- ATF
- ATS
- OpenFOAM (CFD)
- Crunch
- Crunch-Flow
- SWAT
- Machine Learning

File storage and specifications



- Total files per simulation: 5 to a few million files
- Average file size: 100 MB 2 TB
- Total current storage: 100's of MB to 100's of TB (mean = 28 TB/modeler, median = 650 GB/modeler).


Details for typical simulation1 to be archived

Model	Spatial Resolution or Represent- ation	Spatial Extent	Temporal Resolution ²			Mean file size (GB)	Types of file formats	Total annual storage needs (GB)
Multiple LSMs ³	Point ⁴	point	daily	200 yrs	300	0.1	CSV	50
				10 20				
ELM	point	point	hourly, daily	yrs	20	0.004	netCDF	3
ELM	1/2° 2°	global	monthly	250 yrs	2500	0.2	netCDF	15000
ELM-FATES FATES	point, ~1km, ~1degree point	point, regional, and global modes point	sub-daily, monthly <hourly< td=""><td>~500 yrs 10 yrs</td><td>1K 10K 70</td><td>50</td><td>netCDF netCDF</td><td>1000 2000</td></hourly<>	~500 yrs 10 yrs	1K 10K 70	50	netCDF netCDF	1000 2000
		100m 10					HDF5,	
ELM-PFLOTRAN	1 100m	km	hourly/daily	10+ yrs	10 100	10	netCDF	1000
PFLOTRAN	<1m	5-6 km	< hourly	30 yrs	5	1000	HDF5	10000
ATS	100m 250m	10km	daily	10 100 yrs	20	100	XML + HDF5, CSV	1000
ATS	<1 100m	10m 10km	daily	10 100 yrs		2	XML + HDF5	1000
ATS	0.25m	25m	daily	100 yrs	50 200		XML + HDF5	10
CrunchFlow	<1m	<1km	<hourly< td=""><td>30 days</td><td>100</td><td>0.001</td><td>TXT</td><td>1</td></hourly<>	30 days	100	0.001	TXT	1

What model data components are worth archiving?




- Journal and funding requirements for data archiving
- Archive almost everything!
- Exception outputs due to size: archive only high-level outputs corresponding to key figures/findings
- Ambiguity around model code



restart files








Purposes for storing model data publicly



- Transparency and reproducibility
- Model intercomparisons and synthesis
- Comparison with observations
- Standards for ensuring data reusability
 - Machine-readable
 - Critical metadata
 - Workflow documents
 - 90% willing to learn new guidelines or standardized reporting format for model data



Any Questions?







Guidelines: How to archive model data

DATA SCIENCE JOURNAL

Reading: Guidelines for Publicly Archiving Terrestrial Model
Data to Enhance Usability, Intercomparis...

Share: f 💆 🖇 in

Research Papers

Guidelines for Publicly Archiving Terrestrial Model Data to Enhance Usability, Intercomparison, and Synthesis

Authors: Maegen B. Simmonds, William J. Riley, Deborah A. Agarwal, Xingyuan Chen, Shreyas Cholia, Robert Crystal-Ornelas, Ethan T. Coon, Dipankar Dwivedi, Valerie C. Hendrix, Maoyi Huang, Ahmad Jan, Zarine Kakalia, Jitendra Kumar, Charles D. Koven, Li Li, Mario Melara, Lavanya Ramakrishnan, Daniel M. Ricciuto, Anthony P. Walker, Wei Zhi, Oing Zhu, Charuleka Varadharaian

■

Abstract

Scientific communities are increasingly publishing data to evaluate, accredit, and build on published research. However, guidelines for curating data for publication are sparse for model-related research, limiting the usability of archived simulation data. In particular, there are no established guidelines for archiving data related to terrestrial models that simulate land processes and their coupled interactions with climate. Terrestrial modelers have a unique set of challenges when publishing data due to the diversity of scientific domains, research questions, and the types and scales of simulations. Researchers in the U.S. Department of Energy's (DOE) projects use a variety of multiscale models to advance robust predictions of terrestrial and subsurface ecosystem processes. Here, we synthesize archiving needs for data associated with different DOE models, and provide guidelines for publishing terrestrial model data components following FAIR (Findable, Accessible, Interoperable, Reusable) principles. The guidelines recommend archiving model inputs and testing data used in final simulation runs along with associated codes, workflow

Components of model data



- Metadata
- Required Data Files
 - Model Inputs
 - Model outputs
 - Model code
 - Scripts
- Optional Files
 - File-level metadata (FLMD)
 - Model Testing/Validation Data
 - Documentation or user guide

Model Input Files



- Input files required unless publicly available elsewhere
 - Examples: climate forcings, meshes, soil parameterizations
 - Use open-sourced formats such as comma separated value (.csv) or NetCDF (.nc) formats where possible
 - File names should be unique, should only contain letters, numbers, hyphens, underscores, should not contain spaces, and should not rely on case-sensitive file systems
 - Hyperlink to specific input files in metadata and user guide

Use external linking feature if input files are publicly available on another established repository

Model output files



- Includes raw and post-processed data, data supporting findings, tables, figures in a paper.
- Archive all model outputs if the size of the data files are within the repository storage limitations (500GB/file on ESS-DIVE)
- Use decision tree if size of model output files exceed repository storage limits
- Open-sourced formats such as comma separated value (.csv) or NetCDF (.nc) formats where possible.

Contact <u>ess-dive-support@lbl.gov</u> if you have datasets > 0.5 TB






Model Code

- Include source code(s) unless publicly available
- Include specific version, commit hash, or citation allowing exact source code to be recovered
- If available, include external link to tagged release in an established data repository.
- Include links to model codes in metadata/user guide

Scripts

- Include run scripts if necessary for running model to generate results
- Optional: Scripts for reproducing parameters and model configuration for simulations and input files, post-processing model outputs, for executing entire workflow.



Codes published within an ESS-DIVE dataset are CC0 or CCBy4 license. Use external linking feature with software on DOF CODE

Optional Elements



- File-Level Metadata
- Model Testing/Validation Data
 - Data files of observations from each location in an open source format (e.g., CSV).
 - If data are publicly available in another repository, use external linking and include a reference (with DOI) in metadata and user guide.
- Documentation or user guide
 - Readme file (e.g., pdf) for each site-specific or large-scale simulation
 - Provide details on model name, version number, and required data or code dependencies.
 - Include a citation for the model code and licensing information if applicable.

File-level metadata



What are file-level metadata?

 Granular information at the data file level (e.g., file name & description, start and end dates)

Why provide file-level metadata?

- Data users will have general understanding of info contained within a file
- FLMD can enable automatic parsing of data files so that users can eventually search & locate files across data collections



Terri Velliquette

OAK RIDGE

National Laboratory

File-level metadata example



	^	P	С	D
1	File_Name	File_Description	Standard	UTC_offset
2	soil_samples_*.csv	15 soil samples taken in the summer of 2019 using small hand trowel and soil probe.	csv v1.0	- 5 hours
3	SoilPoreWaterHillslope2019.csv	50 soil pore water samples taken from the hillslope at the site over a one year period.	ЕРА	- 5 hours



 FLMD template: https://ess-dive.gitbook.io/file-level-metadata-reporting-format/




Can use wildcard * to indicate when FLMD applies to multiple files

Data package guidelines

Key Criteria

- Authorship
- File size/Repository storage capacity
- Decide what model outputs are worth archiving



Simmonds et al., 2022, Data Sci. Journ.





 Cite dataset and include links to the data and code publication(s) in the Data or Code Availability section and references.

Renteria L Consortiur and Aerob	; Resch C T m T (2020): W ic Respiration or Dynamic Ri	; Tfaily M ; Tolic N ; T PHONDRS Summer 2 I. River Corridor and ver Systems (WHON	y R A; Fansler S; Gara- orgeson J M; Toyoda J 2019 Sampling Campaig Watershed Biogeochem IDRS). doi:10.15485/1729	G; Wells J; Wrighton: Gin: Global River Corristry SFA Worldwide Gin: Global River Corristry SFA Worldwide Gin: Gin: Gin: Gin: Gin: Gin: Gin: Gin:	on K C ; Stegen idor Sediment F Hydrobiogeoch	J C ; WHONDE	RS DC,
		File	s in this dataset Package: ess-	dive-915b1859fbd5ec6-20210	430T040533424808		
► Name	е				File type	Size	Login to Download
	NDRS_Summe	r_2019_Sampling_Camp C_and_Aerobic_Respirat	aign_Global_River_Corridor on.xml	_Sedime	EML v2.1.1	64 KB	Download
⊞ WHO	NDRS_S19S_S	itePhotos.zip		More info	ZIP file	2 GB	Download
⊞ WHO	NDRS_S19S_S	ediment_v2.zip		More info	ZIP file	68 MB	Download
General	Identifier Abstract	The WHONDRS Summand September 2019. 2 downstream). Surface sediment samples and dive.lbl.gov/view/doi:10 surface water and sediscore/transient metabolc This dataset contains (cyclotron resonance m thtps://www.pnnl.gov/er calculated from laborate	high resolution characteriz ass spectrometry (FTICR-MS wironmental-molecular-scier	ampaign collected sample were collected at three warstream site. This data from the surface water. Sasets from this study will as designed with the scibiogeochemical function ation of dissolved organi 5) through the Environme cose-laboratory); (2) non-ss; and (5) metadata. Th	locations within ea set includes a port 519S surface water include geochemic ence community to and physical prop c matter from sedir intal Molecular Scie purgeable organic e field metadata file	ich site (upstream, ion of the data typ r data can be foun cal, hydrologic, and ask questions ass erties of watershe ment via 12 Tesla I ences Laboratory (carbon (NPOC); (e (WHONDRS_S1	midstream, and se produced from the d at https://data.ess-t microbial data from the ociated with links among d and river corridor systems. Fourier transform ion EMSL; 3) respiration rates 95_metadata_v3.esv) is the



Example 1: Citing multiple datasets

Walker, AP, et al. 2019. 'Decadal biomass increment in early secondary succession woody ecosystems is increased by CO2 enrichment'. Nature Communications, 10(1): p. 454. DOI: https://doi.org/10.1038/s41467-019-08348-1

The site-based meteorological dataset

(https://data.ess-dive.lbl.gov/view/ess-dive-7807cf86f1dd42a-20181127T173047368940), the **model output dataset**

(https://data.ess-dive.lbl.gov/view/ess-dive-8260043c35fc925-20181130T171955541030) and the **experiment dataset**

(https://data.ess-dive.lbl.gov/view/ess-dive-f525c71da7d2681-20181128T160851574946) generated and analyzed during the current study are available at the US Department of Energy's (DOE) ESS-DIVE repository.





Koven, CD, et al. 2020. 'Benchmarking and parameter sensitivity of physiological and vegetation dynamics using the Functionally Assembled Terrestrial Ecosystem Simulator (FATES) at Barro Colorado Island, Panama'. Biogeosciences, 17(11): 3017–3044. DOI: https://doi.org/10.5194/bg-17-3017-2020

The **FATES** model is available at https://github.com/NGEET/fates (last access: 15 May 2020; https://doi.org/10.5281/zenodo.3825474, FATES Development Team, 2020). Experiments here are based on git commit 0bc7a5d on the fork: https://github.com/ckoven/fates (last access: 4 June 2020; https://doi.org/10.5281/zenodo.3875687, FATES Development Team, 2019). FATES is run here within two host land surface models, CLM5 and ELMv1, available at https://github.com/ESCOMP/ctsm (git commit b9c92b7, last access: 15 May 2020; https://doi.org/10.5281/zenodo.3739617, CTSM Development Team, 2020) and https://github.com/E3SM-Project/E3SM (git commit 544db3b, last access: 15 May 2020; https://doi.org/10.11578/E3SM/dc.20180418.36, E3SM Project, 2018), respectively. Scripts to initialize parameter files and analyze model output shown here are available at https://github.com/NGEET/testbeds (last access: 15 May 2020; https://doi.org/10.5281/zenodo.3785705, Koven, 2020a), and scripts to run the all model experiments here are available at https://github.com/ckoven/runscripts (last access: 15 May 2020; https://doi.org/10.5281/zenodo.3785703, Koven, 2020b).



Example 3: Citing data, code, scripts

Coon, ET, et al. 2020. 'Coupling surface flow and subsurface flow in complex soil structures using mimetic finite differences'. Advances in Water Resources, 144: 103701. DOI: https://doi.org/10.1016/j.advwatres.2020.103701

The Advanced Terrestrial Simulator (ATS) (Coon et al., 2019) is open source under the BSD 3-clause license and is publicly available at https://github.com/amanzi/ats (last access: October 2019; Coon, 2016). Simulations were conducted using version 0.88. The ATS version 0.88 is permanently stored at https://doi.org/10.5281/zenodo.3727209 (Coon et al., 2020). Forcing data, model input files, Jupyter **notebooks** used to generate figures, and meshes along with Jupyter notebooks used to generate the meshes are publicly available at https://doi.org/10.5440/1545603 (Jan et al., 2019). Data products used in the model comparisons are publicly available through the NGEE Arctic long-term data archive https://doi.org/10.5440/1416559. The observed water level can be accessed at https://doi.org/10.5440/1183767 (Liljedahl and Wilson., 2016), the soil temperature data at https://doi.org/10.5440/1126515 (Romanovsky et al., 2017), and the evapotranspiration data at https://doi.org/10.5440/1362279 (Dengel et al., 2019, respectively.

Any Questions?







ESS-DIVE features and plans to support model data



Application Programming Interface (API) Upload

Web UI limited to 10GB/file, API allows 500GB/file





"External Linking" Feature


Data stored on other repositories are directly linked to dataset metadata on ESS-DIVE and some/all data do not have to be on ESS-DIVE


			aher K; Wainwright H; Williams K (2020): NEON AOP Imaging Spectroscopy sitory. Dataset. doi:el-usecase-1.2 accessed via https://data-dev.ess-dive.lbl.go			Space Radiance and
						Copy Citation
			Files in this dataset Package: ess-dive-64a126fcda197ec-20210930T145535990704			
b	Name			File type	Size	Download All 🕰
B	Metadata: N	EON_AOP_Imaging_Spectroscopy_Survey_of_Upper_East_Ri	ver_Colorado_Watersheds_Raw_Space_Radiance_and_Observational_Variable_Dataset.xml	EML v2.2.0	11 KB	Download 🚨
Externa	I Links to I	ata or Metadata				
			External links for this dataset			
Descripti	on	Relationship	URL			
Flight Dat	a	[has part] Data that is part of this dataset.	https://portal.nersc.gov/wfsfa/doi-10-1	5485-1617204/		
General		the Topical Leaders of				
	Identifier	ess-dive-76578754569b1d4-20210930T145533613560				



Supporting Growth In Data

- Increasing need to support large files and datasets
- Enabling multiple ways to make large data uploads seamless
 - ESS-DIVE API
 - Native ESS-DIVE Uploads
 - Globus
- Upcoming: Tier 2 Storage layer to for very large datasets
 - Supports very large datasets and files
 - Hierarchical data and folders
 - Direct web UI access to browse and download from Tier 2 storage
 - Globus for High Performance Downloads





Globus interface for large scale managed transfers



Upcoming: Tier 2 Storage and Large Uploads



Individual Files over 500GB

Datasets containing any file over 500GB, such as LIDAR or drone data



Over 100 files outside of Zip file

Datasets containing over 100 files that are not stored in a compressed (or "zipped") hierarchy should be treated as large data.

Tier 2 will support uploading large volumes of data; large numbers of files; nested folder structures.

Globus provides high performance interface for data uploads







- ESS-DIVE tracks all versions of data packages internally before and after publication
- Large model data may present challenges with making copies of files with different versions
- Citations indicate version used:

Creator (Publication Year). Title. Publisher. Dataset. Identifier "accessed via data.ess-dive.lbl.gov on YYYY-MM-DD"

Any Questions?







Feedback Tell us what you need!



What are your model data archiving needs?





ESS Modeling Community Needs Summary

- Data are extremely heterogeneous and increasing in volume and complexity
- Disconnect between model and observational data
- Workflows involving manual retrieval of data are not scalable
- Short-term need for most researchers: guidelines to archive model data e.g. those associated with publications
- Long-term: Development of a standardized *model-to-archive* pipeline and *data-to-model* pipeline.





Model-to-archive pipeline

- Community-informed guidelines on creating standardized model data packages
- Pathway for model data packages >500 GB size threshold
- Ability to extract specific subsets of model simulations
- Project portals for sharing and collaborating on pre-published model data
- (long-term) Automating the writing and/or organization of files comprising data packages for specific models or journals.

Data-to-model pipeline

- Support for data formats typically used in model simulations (e.g., netCDF)
- Interoperability between individual data packages in ESS-DIVE and other data centers for model-data integration (field observations and remote sensing data to use for model development, parameterization, and performance testing to improve future measurement designs)

Summary




ESS-DIVE stores model data and has new terrestrial model archiving guidelines

- Model data has many components inputs, outputs, code, documentation
- Publishing model data is beneficial and enables reuse of simulation data
- Guidelines what to archive and how to split datasets for diverse terrestrial model data used by ESS researchers
- ESS-DIVE has many features supporting model data archiving




You can find all this material on ESS-DIVE's...



Model Data Archiving Documentation page

https://docs.ess-dive.lbl.gov/



Webinar page

where this slide deck is available for download

https://ess-dive.lbl.gov/

Additional Resources



Docs: https://ess-dive.gitbook.io/model-data-archiving-guidelines/

Paper: https://datascience.codata.org/articles/10.5334/dsj-2022-003/

Dataset: https://data.ess-dive.lbl.gov/view/doi:10.15485/1813868

Github: https://github.com/ess-dive-community/essdive-model-data-archiving-guidelines

Thank You!





Join ESS-DIVE's Community Mailing List!

http://bit.ly/essdiveMailingList

Contact us at ess-dive-support@lbl.gov